

Evaluation of Exchange-Correlation Energy, Potential, and Stress

L. C. Balbás

Departamento de Física Teórica, Universidad de Valladolid, 47011 Valladolid, Spain.

José Luís Martins

*Departamento de Física, Instituto Superior Técnico, Av Rovisco Pais, 1049-001 Lisboa, Portugal, and
INESC, Rua Alves Redol 9, Apartado 13069, 1000 Lisboa, Portugal.*

José M. Soler

*Departamento de Física de la Materia Condensada, Universidad Autónoma de Madrid, 28049 Madrid, Spain.
(February 1, 2008)*

We describe a method for calculating the exchange and correlation (XC) contributions to the total energy, effective potential, and stress tensor in the generalized gradient approximation. We avoid using the analytical expressions for the functional derivatives of $E_{xc}(\rho)$, which depend on discontinuous second-order derivatives of the electron density ρ . Instead, we first approximate E_{xc} by its integral in a real space grid, and then we evaluate its partial derivatives with respect to the density at the grid points. This ensures the exact consistency between the calculated total energy, potential, and stress, and it avoids the need of second-order derivatives. We show a few applications of the method, which requires only the value of the (spin) electron density in a grid (possibly nonuniform) and returns a conventional (local) XC potential.

PACS:71.15.-m, 71.15.Mb

I. INTRODUCTION

The generalized gradient approximation (GGA)¹ to density functional theory (DFT)² has been growing in acceptance in recent years, due to the development of improved functionals and to the realization of its higher accuracy, for many systems and properties, than that of the local density approximation (LDA)³. Some continuity problems of earlier functionals^{4,5} have been solved in more recent ones⁶. Still, a basic problem is that, while $E_{xc}^{GGA}[\rho]$ depends only on the local value of $\rho(\mathbf{r})$ and $|\nabla\rho(\mathbf{r})|$, its functional derivative depends also on $\nabla|\nabla\rho(\mathbf{r})|$, which is discontinuous where $\nabla\rho(\mathbf{r}) = 0$. This implies that, to avoid aliasing effects, very fine integration grids are required to evaluate the XC potential $v_{xc}(\mathbf{r})$ and its matrix elements. A different problem occurs with grid-oriented implementations of DFT⁷⁻¹¹, in which the electron density is known only at the grid points, while its gradients must be evaluated using finite differences. In this case, an inconsistency between the energy E_{xc} and the potential v_{xc} may occur because different formulas need to be used for the higher-order derivatives of the density. Hamann⁸ proposed an elegant solution to these problems by defining a nonlocal potential, which operates on the *gradient* of the electron wave functions. Although in principle this does not pose any special difficulty, it requires a specific, unconventional implementation, which may be difficult to adapt to existing DFT codes. White and Bird¹² found another solution, by *defining* E_{xc}^{GGA} as an integral in a real space grid (what is always done in practice), and $v_{xc}(\mathbf{r}_i)$ as its *partial* derivative (as opposed to functional derivative) with respect to the density ρ_i at the grid points $\mathbf{r}_i$. They applied this method to a plane wave basis set and a uniform integration grid, finding the

gradients with fast Fourier transforms (FFTs). Here we generalize their method to arbitrary bases and nonuniform grids, and we calculate the density gradient using finite differences. The method produces a standard, local potential, which is exactly consistent with the definition of E_{xc} , in the sense that v_{xc} is the correct potential in the Schrödinger equation that results from the variational minimization of the total energy. In addition, we show how to evaluate the XC contribution to the stress tensor in crystals, following the same ideas. The trivial particular case of the LDA is also discussed.

II. EXCHANGE-CORRELATION ENERGY AND POTENTIAL

The LDA³ and GGA¹ approximations to E_{xc} have the general forms

$$E_{xc}^{LDA}[\rho] = \int f_{LDA}(\rho(\mathbf{r})) d\mathbf{r},$$

$$E_{xc}^{GGA}[\rho] = \int f_{GGA}(\rho(\mathbf{r}), \mathbf{g}(\mathbf{r})) d\mathbf{r}, \quad (1)$$

where $f = \rho \epsilon_{xc}$ is the local XC energy density, and ϵ_{xc} is the XC energy per electron. We use the notation¹³ $\mathbf{g}(\mathbf{r}) \equiv \nabla\rho(\mathbf{r})$, $g_\alpha(\mathbf{r}) \equiv \nabla_\alpha\rho(\mathbf{r})$, and $g(\mathbf{r}) \equiv |\nabla\rho(\mathbf{r})|$. The argument $\mathbf{g}(\mathbf{r})$ in Eq.(1), rather than $g(\mathbf{r})$, indicates that, in principle, f_{GGA} might depend on the relative orientation of the gradients of the two spin components of the density. However, for notational simplicity we will omit the spin index and the sum over it, since their inclusion is trivial. In the limit of slowly varying electron density, we must recover the LDA result, that is $f_{GGA}(\rho, 0) = f_{LDA}(\rho)$ ⁵.

In the LDA the XC potential has a simple expression,

$$v_{\text{xc}}^{\text{LDA}} = \frac{df_{\text{LDA}}}{d\rho} = \epsilon_{\text{xc}}^{\text{LDA}} + \rho \frac{d\epsilon_{\text{xc}}^{\text{LDA}}}{d\rho},$$

whereas in the GGA it is considerably more complicated

$$\begin{aligned} v_{\text{xc}}^{\text{GGA}} &= \frac{\partial f_{\text{GGA}}}{\partial \rho} - \nabla \frac{\partial f_{\text{GGA}}}{\partial \mathbf{g}} \\ &= \epsilon_{\text{xc}}^{\text{GGA}} + \rho \frac{\partial \epsilon_{\text{xc}}^{\text{GGA}}}{\partial \rho} - g \frac{\partial \epsilon_{\text{xc}}^{\text{GGA}}}{\partial g} - \rho g \frac{\partial^2 \epsilon_{\text{xc}}^{\text{GGA}}}{\partial g \partial \rho} \\ &\quad - \frac{\rho}{g} \frac{\partial \epsilon_{\text{xc}}^{\text{GGA}}}{\partial g} \nabla^2 \rho + \frac{\rho}{g^2} \frac{\partial \epsilon_{\text{xc}}^{\text{GGA}}}{\partial g} \mathbf{g} \nabla g \\ &\quad - \frac{\rho}{g} \frac{\partial^2 \epsilon_{\text{xc}}^{\text{GGA}}}{\partial g^2} \mathbf{g} \nabla g, \end{aligned} \quad (2)$$

as it depends on the first and second gradients of the electron density. Since $|\nabla \rho|$ has cusps at extrema of ρ , ∇g is discontinuous at these points, what causes problems for its numerical representation. Furthermore, some parameterizations of $\epsilon_{\text{xc}}^{\text{GGA}}$ which apparently join seamlessly different density regimes, may have higher derivatives which do not behave well in those transition regions.

In what follows, we will omit the label GGA, except to underline a distinction with LDA. In a practical calculation, the XC energy is calculated through a numerical integration. From a set of M mesh points $\mathbf{r}_i$ and weights w_i , we approximate

$$E_{\text{xc}} \simeq \sum_{i=1}^M w_i f(\rho(\mathbf{r}_i), \mathbf{g}(\mathbf{r}_i)).$$

The weights may be, for example $4\pi r_i^2 \Delta r$ in a uniform radial grid, or the jacobian of the local metric tensor in an adaptive-coordinate grid¹⁴ (see below). In addition, we must specify precisely the meaning of $\mathbf{g}(\mathbf{r}_i)$. If a well defined basis set is used, the electron density and its gradient can be calculated exactly at any point in space, from the electron wave functions and their gradients. In practice, this may add an appreciable overhead in terms of computer time and memory. Alternatively, we can use the values of the density at the grid points to calculate its gradient, using either FFTs¹² or finite differences. As the gradient is a linear operator, we can write in general

$$g_{\alpha}(\mathbf{r}_i) \simeq g_{i\alpha} \equiv \sum_{j=1}^M A_{ij}^{\alpha} \rho_j, \quad (3)$$

and $\partial g_{i\alpha} / \partial \rho_j = A_{ij}^{\alpha}$. $\{i, j\}$ are combined indexes ($i \equiv \{i_1, i_2, i_3\}$) that label grid points, α labels the three cartesian coordinates, $\rho_i \equiv \rho(\mathbf{r}_i)$, and the coefficients A_{ij}^{α} depend on the mesh and the chosen numerical derivative formula, but not on ρ_i . There are many choices for the coefficients A_{ij}^{α} , depending on the integration mesh and interpolation method, since the numerical derivative can be defined as the derivative of the interpolation function.

The particular case described by White and Bird¹² uses a uniform mesh and a Fourier-series interpolation. It was developed for plane-wave calculations, and FFTs were used to evaluate Eq. (3). Our method can be generalized to nonuniform grids. In the implementation that we will describe later, we use a local Lagrange interpolation, for which the matrix A_{ij}^{α} is sparse.

Notice that in Eq. (3) we are *defining* $\mathbf{g}_i$ as the numerical derivative of ρ on the mesh (therefore being a function of the values $\{\rho_j\}$), while we reserve the notation $\mathbf{g}(\mathbf{r}_i)$ for the exact gradient of ρ at $\mathbf{r}_i$ (in case it is known). It may be argued that the use of $\mathbf{g}_i$, instead of $\mathbf{g}(\mathbf{r}_i)$ represents an additional approximation. However, in practice $\mathbf{g}_i$ is frequently a better approximation than $\mathbf{g}(\mathbf{r}_i)$ to the *average* value of $\mathbf{g}(\mathbf{r})$ within the spatial “pixel” which corresponds to the integration point $\mathbf{r}_i$. Both values agree for a Fourier interpolation, provided that the plane wave cutoff of the grid is twice as large as that of the wave functions. And of course they must also agree, for any interpolation scheme, in the limit of an infinitely fine grid.

Thus, following White and Bird¹², we *define*

$$\tilde{E}_{\text{xc}}(\{\rho_i\}) \equiv \sum_{i=1}^M w_i f_i(\rho_i, \mathbf{g}_i(\{\rho_j\})) \simeq E_{\text{xc}}, \quad (4)$$

and

$$\begin{aligned} \tilde{v}_i^{\text{xc}} &\equiv \frac{1}{w_i} \frac{\partial \tilde{E}_{\text{xc}}(\{\rho_j\})}{\partial \rho_i} \\ &= \frac{\partial f}{\partial \rho_i} + \sum_{j=1}^M \sum_{\alpha=1}^3 \frac{w_j}{w_i} \frac{\partial f_j}{\partial g_{j\alpha}} \frac{\partial g_{j\alpha}}{\partial \rho_i}. \end{aligned} \quad (5)$$

It is important to emphasize that Eqs. (2) and (5) are asymptotically equivalent in the limit of an infinitely fine grid, but different in practice (in particular, Eq. (5) has no explicit dependence on the second derivatives of ρ). In fact, it is easy to see that Eq. (5), and not Eq. (2), is the “correct” definition of v_{xc} , if the functional form (4) is actually used in the variational energy minimization, $\partial E / \partial \psi^* = H\psi$, because, from Eq. (5),

$$\frac{\partial \tilde{E}_{\text{xc}}}{\partial \psi_i^*} = \frac{\partial \tilde{E}_{\text{xc}}}{\partial \rho_i} \frac{\partial \rho_i}{\partial \psi_i^*} = w_i \tilde{v}_i^{\text{xc}} \psi_i.$$

In the LDA, f_i does not depend on $\mathbf{g}_i$, and only the first term in Eq. (5) remains, giving the trivial result

$$\tilde{v}_i^{\text{LDA}} = \frac{df_i^{\text{LDA}}}{d\rho_i}.$$

There remains the problem of determining the coefficients A_{ij}^{α} and the weights w_i in Eqs. (3) and (4). To this purpose, it is convenient to introduce grid variables $\{s_{\mu}; \mu = 1, 2, 3\}$, which are in principle continuous. In practice, however, the density $\rho(\mathbf{s})$ and the cartesian coordinates $\mathbf{r}(\mathbf{s})$ of the grid points are evaluated only at integer values of s_{μ} . For a regular grid,

$r_\alpha(\mathbf{s}) = \sum_{\mu=1}^3 s_\mu a_{\mu\alpha}/N_\mu$, where $a_{\mu\alpha}$ is the α th cartesian coordinate of the μ th lattice unit vector, and N_μ is the number of grid divisions along that vector. For nonuniform grids, $r_\alpha(\mathbf{s})$ are general functions and it is convenient to introduce the jacobian of the transformation $\mathbf{s} \rightarrow \mathbf{r}$. The XC energy can then be expressed as

$$E_{\text{xc}} = \int f(\rho(\mathbf{s}), \mathbf{g}(\mathbf{s})) \left| \frac{\partial \mathbf{r}}{\partial \mathbf{s}} \right| d\mathbf{s}$$

or

$$\tilde{E}_{\text{xc}} = \sum_{i=1}^M f(\rho_i, \mathbf{g}_i) |D_i|$$

where we have used that $\Delta s_\mu = 1$ by definition of s_μ . $|D_i|$ is the determinant at point i of the matrix of partial derivatives

$$D_{\alpha\mu}(\mathbf{s}) \equiv \frac{\partial r_\alpha(\mathbf{s})}{\partial s_\mu}.$$

By comparison with (4), we conclude that $w_i = |D_i|$. Since the grid variables s_μ are by definition ‘‘orthogonal’’ and the functions r_α and ρ are evaluated at regular unit intervals of s_μ , their derivatives $\partial/\partial s_\mu$ can be calculated straightforwardly:

$$D_i^{\mu\alpha} = \left(\frac{\partial r_\alpha}{\partial s_\mu} \right)_i = \sum_{j=1}^M B_{ij}^\mu r_{j\alpha} \quad (6)$$

$$\left(\frac{\partial \rho}{\partial s_\mu} \right)_i = \sum_{j=1}^M B_{ij}^\mu \rho_j$$

where the coefficients B_{ij}^μ depend only on the relative values of i and j , and are independent of the grid coordinates $\mathbf{r}(\mathbf{s})$:

$$B_{ij}^\mu = \begin{cases} L_k^{(n)} & \text{if } j_\mu = i_\mu + k; j_\nu = i_\nu, \nu \neq \mu \\ -L_k^{(n)} & \text{if } j_\mu = i_\mu - k; j_\nu = i_\nu, \nu \neq \mu \\ 0 & \text{otherwise} \end{cases}$$

where the coefficients $L_k^{(n)}$ may be derived from a $(2n+1)$ -point Lagrange polynomial interpolation formula²⁴.

We can now calculate $\mathbf{g}$ as

$$g_{i\alpha} = \sum_{\mu=1}^3 \left(\frac{\partial \rho}{\partial s_\mu} \right)_i \left(\frac{\partial s_\mu}{\partial r_\alpha} \right)_i = \sum_{\mu=1}^3 \sum_{j=1}^M (D_i^{-1})_{\alpha\mu} B_{ij}^\mu \rho_j$$

and by identifying coefficients with Eq. (3)

$$A_{ji}^\alpha = \frac{\partial g_{j\alpha}}{\partial \rho_i} = \sum_{\mu=1}^3 (D_j^{-1})_{\alpha\mu} B_{ji}^\mu. \quad (7)$$

Thus, the calculation of the XC potential from the density on the grid involves the following steps, for every

point j : 1) find the 3×3 matrix D_j , using Eq. (6), its determinant w_j , and its inverse D_j^{-1} , storing w_j for later use; 2) find A_{ji}^α from Eq. (7) and $\mathbf{g}_j$ from Eq. (3); 3) calculate $f(\rho_j, \mathbf{g}_j)$ and its derivatives with respect to ρ_j and $g_{j\alpha}$; 4) add the first term in Eq. (5) (multiplied by w_j) to $\tilde{v}_j^{\text{xc}}$ and the second term (except the denominator w_i) to $\tilde{v}_i^{\text{xc}}$ for every neighbor point i involved in the calculation of $\mathbf{g}_j$; 5) when the previous loop is finished, run again over all grid points i , dividing $\tilde{v}_i^{\text{xc}}$ by w_i . In the case of a uniform grid, the matrix D_i does not depend on i . Thus, w_i and D_i^{-1} can be evaluated once and for all, saving steps 1 and 5, as well as the temporary array required to store w_i .

III. EXCHANGE-CORRELATION CONTRIBUTION TO THE STRESS TENSOR

We consider now the stress tensor¹⁵

$$\sigma_{\alpha\beta}^{\text{xc}} \equiv \frac{\partial \tilde{E}_{\text{xc}}}{\partial \epsilon_{\alpha\beta}}$$

where $\epsilon_{\alpha\beta}$ is the strain tensor, giving the deformation of all points in space (including atomic and grid coordinates):

$$r_\alpha \rightarrow \sum_{\beta=1}^3 (\delta_{\alpha\beta} + \epsilon_{\alpha\beta}) r_\beta. \quad (8)$$

More generally, we consider the derivative of $\tilde{E}_{\text{xc}}$ with respect to a parameter λ that affects the position of the grid points. This may be $\epsilon_{\alpha\beta}$ or one of the atomic positions, in case of a nonuniform grid which depends on them¹⁴. It is therefore convenient to recognize explicitly that, when the system is modified, $\tilde{E}_{\text{xc}}$ depends on the grid point coordinates in addition to the densities at the grid points, i.e. $\tilde{E}_{\text{xc}}(\{\rho_i\}, \{\mathbf{r}_i\})$. Thus,

$$\frac{\partial \tilde{E}_{\text{xc}}}{\partial \lambda} = \sum_{i=1}^M \left(\frac{\partial \tilde{E}_{\text{xc}}}{\partial \rho_i} \frac{\partial \rho_i}{\partial \lambda} + \sum_{\alpha=1}^3 \frac{\partial \tilde{E}_{\text{xc}}}{\partial r_{i\alpha}} \frac{\partial r_{i\alpha}}{\partial \lambda} \right)$$

where $\partial \tilde{E}_{\text{xc}}/\partial \rho_i = \tilde{v}_i^{\text{xc}}$ and

$$\frac{\partial \tilde{E}_{\text{xc}}}{\partial r_{i\alpha}} = \sum_{j=1}^M \left(\frac{\partial |D_j|}{\partial r_{i\alpha}} f_j + |D_j| \sum_{\beta=1}^3 \frac{\partial f_j}{\partial g_{j\beta}} \frac{\partial g_{j\beta}}{\partial r_{i\alpha}} \right)$$

$$\frac{\partial g_{j\beta}}{\partial r_{i\alpha}} = \sum_{\mu=1}^3 \left(\frac{\partial \rho}{\partial s_\mu} \right)_j \frac{\partial}{\partial r_{i\alpha}} (D_j^{-1})_{\beta\mu}$$

Now, using Eq. (8) we find

$$\frac{\partial r_\gamma}{\partial \epsilon_{\alpha\beta}} = \delta_{\gamma\alpha} r_\beta, \quad (9)$$

and taking into account the general properties of a determinant, as well as the fact that $(DD^{-1} = I) \Rightarrow (\delta D = -D^{-1}\delta D D^{-1})$, we finally find

$$\frac{\partial \tilde{E}_{xc}}{\partial \epsilon_{\alpha\beta}} = \delta_{\alpha\beta} \tilde{E}_{xc} + \sum_{i=1}^M w_i \tilde{v}_i^{xc} \frac{\partial \rho_i}{\partial \epsilon_{\alpha\beta}} - \sum_{i=1}^M w_i \frac{\partial f_i}{\partial g_{i\beta}} g_{j\alpha} \quad (10)$$

In the limit of an infinitely fine grid, we have

$$\begin{aligned} \frac{\partial E_{xc}}{\partial \epsilon_{\alpha\beta}} &= \delta_{\alpha\beta} E_{xc} + \int v_{xc}(\mathbf{r}) \frac{\partial \rho(\mathbf{r})}{\partial \epsilon_{\alpha\beta}} d\mathbf{r} \\ &\quad - \int \frac{\partial f(\rho(\mathbf{r}), \mathbf{g}(\mathbf{r}))}{\partial g_{\beta}(\mathbf{r})} g_{\alpha}(\mathbf{r}) d\mathbf{r}. \end{aligned} \quad (11)$$

The presence of $\partial \rho / \epsilon_{\alpha\beta}$ in Eqs. (10) and (11) requires to clarify what is kept constant when taking the derivatives. We will argue that the “correct” definition (or at least the most useful one) requires to keep constant the variational coefficients c_{na} of the expansion of the electron wave functions $\psi_n(\mathbf{r})$ in terms of the N basis functions $\phi_a(\mathbf{r})$:

$$\psi_n(\mathbf{r}) = \sum_{a=1}^N c_{na} \phi_a(\mathbf{r}) \quad (12)$$

Since $\sigma_{\alpha\beta}$ is generally evaluated at the electronic ground state, the Hellman-Feynman theorem ensures that the change of c_{na} will not affect the total energy to first order. Of course, the Hellman-Feynman theorem does not apply to E_{xc} alone, but in practice we are interested in adding $\sigma_{\alpha\beta}^{xc}$ to other contributions of $\sigma_{\alpha\beta}$, in order to calculate the total stress tensor. And even in the Car-Parrinello¹⁹ method, in which forces and stresses are evaluated out of the ground state, the appropriate definition of these magnitudes involves the derivatives of the total energy at constant c_{na} .

There are two reasons why ρ_i depends on $\epsilon_{\alpha\beta}$: the change of the basis functions $\phi_a(\mathbf{r}_i)$ at the displaced grid points, and the change of the wave function coefficients c_{na} required to maintain the orthonormality constraints $\langle \psi_n | \psi_m \rangle = \delta_{nm}$. With a plane wave basis set, or with a grid-oriented scheme^{7,10,9}, in which the wave functions are defined directly at the grid points, without any basis, the orthonormality constraints are not affected by the deformation of the unit cell, provided that the wave functions and the density are simply scaled by a factor $(\Omega_0/\Omega)^{1/2}$ and Ω_0/Ω , respectively, where Ω_0 and Ω are the unit cell volumes before and after the deformation. It is easy to verify that $\partial \Omega / \partial \epsilon_{\alpha\beta} = \Omega \delta_{\alpha\beta}$ and therefore

$$\frac{\partial \rho(\mathbf{r})}{\partial \epsilon_{\alpha\beta}} = -\delta_{\alpha\beta} \rho(\mathbf{r}) \quad (13)$$

Substitution of (13) into (11) leads to

$$\begin{aligned} \frac{\partial E_{xc}}{\partial \epsilon_{\alpha\beta}} &= \delta_{\alpha\beta} \int (\epsilon_{xc}(\mathbf{r}) - v_{xc}(\mathbf{r})) \rho(\mathbf{r}) d\mathbf{r} \\ &\quad - \int \frac{\partial f(\rho(\mathbf{r}), \mathbf{g}(\mathbf{r}))}{\partial g_{\beta}(\mathbf{r})} g_{\alpha}(\mathbf{r}) d\mathbf{r}, \end{aligned}$$

which coincides with Eq. (24) of Dal Corso and Resta¹⁸.

In the case of an atomic basis set, it is convenient to define the density matrix

$$\rho_{ab} = \sum_{n=1}^N q_n c_{na}^* c_{nb}$$

with q_n the occupation of state ψ_n . Then

$$\begin{aligned} \rho(\mathbf{r}) &= \sum_{a,b=1}^N \rho_{ab} \phi_a(\mathbf{r}) \phi_b(\mathbf{r}) \\ \frac{\partial \rho(\mathbf{r})}{\partial \lambda} &= \sum_{a,b=1}^N \left(\frac{\partial \rho_{ab}}{\partial \lambda} \phi_a(\mathbf{r}) \phi_b(\mathbf{r}) \right. \\ &\quad \left. + 2\rho_{ab} \phi_a(\mathbf{r}) \sum_{\alpha=1}^3 \nabla_{\alpha} \phi_b(\mathbf{r}) \frac{\partial (r_{\alpha} - R_{b\alpha})}{\partial \lambda} \right) \end{aligned} \quad (14)$$

where we are assuming real basis orbitals for simplicity. $\mathbf{R}_b$ is the origin (atomic position) of orbital ϕ_b , and the last term in Eq. (14) accounts for the change in the relative position $\mathbf{r} - \mathbf{R}_b$ when we deform the lattice or change the grid point positions (we assume a constant shape of ϕ_b). $\partial \rho_{ab} / \partial \lambda$ is the change in the density matrix required to maintain the orthonormality constraints:

$$\frac{\partial \rho_{ab}}{\partial \lambda} = - \sum_{c=1}^N \rho_{ac} \frac{\partial S_{cb}}{\partial \lambda} = - \sum_{c=1}^N \rho_{ac} \sum_{\alpha=1}^3 \frac{\partial S_{cb}}{\partial r_{cb}^{\alpha}} \frac{\partial r_{cb}^{\alpha}}{\partial \lambda} \quad (15)$$

where $S_{ab} \equiv \langle \phi_a | \phi_b \rangle$, and $\mathbf{r}_{ab} \equiv \mathbf{R}_b - \mathbf{R}_a$. By combining Eqs. (9), (14), and (15) we obtain

$$\begin{aligned} \int v_{xc}(\mathbf{r}) \frac{\partial \rho(\mathbf{r})}{\partial \epsilon_{\alpha\beta}} d\mathbf{r} &= - \sum_{a,b,c=1}^N \rho_{ac} \frac{\partial S_{cb}}{\partial r_{cb}^{\alpha}} r_{cb}^{\beta} \langle \phi_a | v_{xc} | \phi_b \rangle \\ &\quad + 2 \sum_{a,b=1}^N \rho_{ab} \langle \phi_a | v_{xc}(\mathbf{r}) (r_{\beta} - R_{b\beta}) \nabla_{\alpha} | \phi_b \rangle. \end{aligned}$$

The same expression is valid if the integral is replaced by the grid sum $\sum_i w_i \tilde{v}_i^{xc} (\partial \rho_i / \partial \epsilon_{\alpha\beta})$, except that then the matrix elements must also be calculated on the grid, i.e.

$$\langle \phi_a | v_{xc} | \phi_b \rangle = \sum_{i=1}^M w_i \tilde{v}_i^{xc} \phi_{ai} \phi_{bi}$$

Similar terms appear in other contributions to the stress tensor. Thus, in the derivation of the Hartree energy, there is an identical term, except for the substitution of v_{xc} by the Hartree potential v_H . Then the total contribution to the stress tensor that arises from the change in the density has the form

$$\begin{aligned} \int \frac{\delta E}{\delta \rho(\mathbf{r})} \frac{\partial \rho(\mathbf{r})}{\partial \epsilon_{\alpha\beta}} d\mathbf{r} &= - \sum_{a,b=1}^N E_{ab} \frac{\partial S_{ab}}{\partial r_{ab}^{\alpha}} r_{ab}^{\beta} \\ &\quad + 2 \sum_{a,b=1}^N \rho_{ab} \langle \phi_a | v(\mathbf{r}) (r_{\beta} - R_{b\beta}) \nabla_{\alpha} | \phi_b \rangle \end{aligned} \quad (16)$$

where $v(\mathbf{r})$ is the total effective potential, and E_{ab} are energy-density matrix elements,

$$E_{ab} = \sum_{c=1}^N \rho_{ac} H_{cb} = \sum_{n=1}^N q_n \epsilon_n c_{na}^* c_{nb}, \quad (17)$$

with $H_{ab} = \langle \phi_a | H | \phi_b \rangle$, H the total one-electron hamiltonian and ϵ_n its n th eigenvalue. Notice, however, that the derivation of Eqs. (16) and (17) does *not* assume that the matrix elements S_{ab} and H_{ab} or their derivatives are calculated on the grid. In fact, in our atomic-basis DFT implementation²⁰, two-center matrix elements like S_{ab} or $\langle \phi_a | -\nabla^2 | \phi_b \rangle$, are calculated using reciprocal-space convolution techniques²¹.

IV. ACCURACY TESTS

We have implemented the method described above as an independent package, which we are using with several electronic structure programs, including an atomic pseudopotential generator, the localized-basis program SIESTA²⁰ and a pseudopotential plane-wave program.

We have implemented two different LDA parameterizations, PZ-LDA²² and PW92-LDA⁴, of Ceperley and Adler electron gas energies²³ for f_{LDA} and two different GGA recipes, PW91-GGA⁴ and PBE-GGA⁶ for f_{GGA} . They were all implemented in their spin-dependent formulations. As the calculation of f and its derivatives are kept as separate procedures it is easy to add new parameterizations. As only first derivatives of f are needed in our method to calculate stresses and potentials, we do not have to calculate second derivatives as in the traditional method. Since GGA functional forms tend to be complicated, this is convenient.

In the atomic pseudopotential generation program the electron density is assumed to have spherical symmetry. The radial mesh points are derived from a monotonous function $r(s)$. The gradient of the density is calculated analogously to the three-dimensional case:

$$g_i = \left(\frac{dr}{ds} \right)_i^{-1} \left(\frac{d\rho}{ds} \right)_i,$$

$$\left(\frac{dr}{ds} \right)_i = \sum_{k=1}^n L_k^{(n)} (r_{i+k} - r_{i-k}),$$

$$\left(\frac{d\rho}{ds} \right)_i = \sum_{k=1}^n L_k^{(n)} (\rho_{i+k} - \rho_{i-k}).$$

To illustrate the method we have chosen three simple situations. Fig. 1 shows the s -pseudopotential for Si, with the Troullier-Martins recipe²⁵ and a core radius of two bohr, calculated with the PBE-GGA and PW91-GGA functionals using both our method (Eq. (5)) and




FIG. 1. s -pseudopotential generated for Si. The solid curve was generated with the PW91-GGA and the dashed curve with the PBE-GGA. The curves generated with the traditional and the new methods are undistinguishable.

the usual method (Eq. (2)), with the standard radial grid used by the pseudopotential generation program. The curves for the two computational methods agree within 10^{-4} Ry and are indistinguishable in the scale of the figure. The PW91-GGA pseudopotential has some wiggles that are due to the term proportional to $\exp(-100s^2)$ where $s = \nabla\rho/(2k_F\rho)$, $k_F = (3\pi^2\rho)^{1/3}$, in the parameterization of the GGA exchange. Those wiggles occur near the extrema of the valence electron density, and they are not present in the PBE-GGA pseudopotential, which uses a different parameterization of the GGA exchange. By construction the screened pseudopotential is smooth, without wiggles, so the appearance of those wiggles are due to the unscreening of the pseudopotential. To see any difference between the usual and present methods, we must use very coarse radial meshes. This is shown in Fig. 2, where we compare the XC potential for a model density $\rho(\mathbf{r}) = (\sin r/r)^2$. The PZ-LDA (dotted thin line), PW91-GGA (solid thick line), and PBE-GGA (dashed line), were first calculated on a fine radial mesh. Again there is no difference between the old and new methods on the scale of fig. 2 for the fine mesh. However, recalculating the PW91-GGA exchange and correlation potential on a much coarser mesh with the new method (squares) one starts to see small differences in the potential values in the regions where the PW91 parameterization has wiggles. The eleven point Lagrange formula for coarse sampling spans to a region of 2 bohr, almost half of the horizontal range of Fig. 2. It misses the small wiggles around $r = 3\pi/2$ where $\nabla\rho = 0$ and $\rho \neq 0$, but it finds wiggles at $r = \pi$ where both $\nabla\rho = 0$ and $\rho = 0$.

Our last example uses the pseudo-charge density of diamond. A well converged plane wave expansion of the density is used to calculate ρ_i for different $N \times N \times N$ grids. Fig. 3 shows that the calculated XC energy is




FIG. 2. XC potential for a spherical electron density with the form $\rho(r) = (\sin r/r)^2$. The dashed line is the CA-LDA, the dotted line is the PBE-GGA and the solid line is the PW91-GGA, all accurately calculated on a fine grid with a step of 0.02 bohr. The squares show that the PW91-GGA, calculated with the new method on a coarse grid (step = 0.2 bohr) have small deviations with respect to the fine grid. However, in fact these deviations tend to smooth out the large pathological wiggles developed by the PW91-GGA functional in regions of zero density or zero density gradient. This makes the present method very well behaved even with very coarse grids.

very stable with respect to grid size: even a $8 \times 8 \times 8$ grid gives a value accurate within 10^{-3} Hartree. The XC component of the stress also converges very fast, and the method is very stable. In practice, the PW cutoffs required for a good convergence of the total energy impose larger grids than those needed to converge E_{xc} with our method. Thus, the standard grids used in typical PW calculations are more than sufficient for E_{xc} .

V. CONCLUSIONS

We have derived formulas for the GGA-XC energy, potential, and stress, as a function of the electron density given in a spatial mesh of points, which may be unevenly distributed. Density gradients need not be provided, since they are calculated numerically using the density at the grid points, what leads to well behaved formulas. The use of a unique definition of the gradients ensures an *exact consistency* between the calculated values for the energy, potential, and stress, including all corrections for basis set incompleteness and changing grid points. As the number of grid points increases, we recover the exact results obtained using the virial theorem.




FIG. 3. XC energy in diamond as a function of the grid size $N \times N \times N$. Squares are from LDA and circles from GGA. The charge density $\rho(r)$ is the same in all cases. Only the sampling grid and the XC functional change.

ACKNOWLEDGMENTS

This work was supported by the Fundación Ramón Areces and by MCT/DGI grants PB00-1312 and PB98-0368-C02-01.

- ¹ D. C. Langreth and M. J. Mehl, Phys. Rev. B **28**, 1809 (1983). A. D. Becke, Phys. Rev. A, **38**, 3098 (1988).
- ² P. Hohenberg and W. Kohn, Phys. Rev. **136**, B864 (1964).
- ³ W. Kohn and L. J. Sham, Phys. Rev. **136**, A1133 (1965).
- ⁴ J. P. Perdew and Y. Wang, Phys. Rev. B **45**, 13244 (1992). J. P. Perdew in *Electronic Structure of Solids 91*, p. 11, edited by P. Ziesche and H. Eschrig (Akademie Verlag, Berlin, 1991).
- ⁵ C. Filippi, C. J. Umrigar and M. Taut, J. Chem. Phys. **100**, 1290 (1994). C. Filippi, X. Gonze and C. J. Umrigar in *Recent Developments and Applications of Modern Density Functional Theory*, p. 295, edited by J. M. Seminario (Elsevier, 1996).
- ⁶ J. P. Perdew, K. Burke, and M. Ernzerhof, Phys. Rev. Lett. **77**, 3865 (1996). We are not considering a new generation of GGA functionals including a dependence with the laplacian of the electron density. Nevertheless, the main approach in this paper is even more useful for that type of functionals. See, i.e., A. Rassolov, J. A. Pople and M. A. Ratner, Phys. Rev. B **59**, 15625 (1999); *ibid* **62**, 2232 (2000).
- ⁷ J. R. Chelikowsky, N. Troullier, and Y. Saad, Phys. Rev. Lett. **72**, 1240 (1994); J. R. Chelikowsky, N. Troullier, K. Wu, and Y. Saad, Phys. Rev. B **50**, 11355 (1994).
- ⁸ D. R. Hamann, Phys. Rev. B **54**, 1568 (1996).
- ⁹ N. A. Modine, G. Zumbach, and E. Kaxiras, Phys. Rev. B **55**, 10289 (1997).

- ¹⁰ E. L. Briggs, D. J. Sullivan, and J. Bernholc, Phys. Rev. B **52**, R5471 (1995); *ibid* **54**, 14362 (1996).
- ¹¹ T. L. Beck, Rev. Modern Phys. **72**, 1041 (2000)
- ¹² M.J. A. White and D. M. Bird, Phys. Rev. B **50**, 4954 (1994).
- ¹³ For convenience, we will place indexes and labels indistinctly as sub or superscripts, without meaning a tensorial notation nor implicit summations. Indexes i, j are used for the M mesh points, α, β, γ for the three cartesian coordinates, μ, ν for the three lattice or mesh coordinates, a, b, c for the N basis functions, and n, m for the N hamiltonian eigenstates.
- ¹⁴ F. Gygi, Phys. Rev. B **48**, 11692 (1993); F. Gygi and G. Galli, Phys. Rev. B **52**, R2229 (1995).
- ¹⁵ We use here the convention that the stress has the opposite sign than the pressure. Often the opposite convention is used, so the reader may find an overall change of sign in the expressions given by different authors.
- ¹⁶ As the energy of a crystal does not depend on its orientation in space, only the length of the lattice vectors and the angles between them are important. In other words, the metric tensor $G_{\mu\nu} = \mathbf{a}_\mu \cdot \mathbf{a}_\nu$ contains all the relevant information. This allows an equivalent formulation, in which the contravariant components of the stress tensor can be defined as $\tau^{\mu\nu} = 2\partial E / \partial G_{\mu\nu}$ and its cartesian components are then $\sigma_{\alpha\beta} = \sum_{\mu\nu} \tau^{\mu\nu} a_{\mu\alpha} a_{\nu\beta}$. For details, see I. Souza and J. L. Martins, Phys. Rev. B **55**, 8733 (1997).
- ¹⁷ O. H. Nielsen and R. M. Martin, Phys. Rev. B **32**, 3780 (1985).
- ¹⁸ A. Dal Corso and R. Resta, Phys. Rev. B **50**, 4327 (1994).
- ¹⁹ R. Car and M. Parrinello, Phys. Rev. Lett. **55**, 2471 (1985).
- ²⁰ P. Ordejón, E. Artacho, and J. M. Soler, Phys. Rev. B **53**, R10441 (1996); D. Sánchez-Portal, P. Ordejón, E. Artacho, and J. M. Soler, Int. J. Quant. Chem. **65**, 453 (1997).
- ²¹ O. F. Sankey and D. J. Niklewski, Phys. Rev. B **40**, 3979 (1989).
- ²² J. P. Perdew and A. Zunger, Phys. Rev. B **23**, 5048 (1981).
- ²³ D. M. Ceperley and B. J. Alder, Phys. Rev. Lett. **45**, 566 (1980).
- ²⁴ *Handbook of Mathematical Tables*, Ed. by M. Abramowitz and I. A. Stegun (Dover, New York, 1965).
- ²⁵ N. Troullier and J. L. Martins, Phys. Rev. B **43**, 1993 (1991).